

Phytoremediation of salt-affected soils in the Toronto region

Lauren Nawroth, MES Candidate
Queen's University

Lyndsay Cartwright, PhD
Senior Research Analyst, Toronto and Region Conservation Authority

Barb Zeeb, PhD
Professor and Canada Research Chair in Biotechnology and the Environment
Royal Military College

Regional Watershed Alliance

September 18, 2024



Acknowledgements



TERRERA










Chloride levels in freshwater continue to rise





Lake Wilcox





Well ID 382-1 (Lower Newmarket formation)



Moving towards, or already exceeding, thresholds to protect aquatic life & human well-being



Legacy Chloride



Legacy Chloride



Salinity Stress



Healthy Spreading Orache leaf¹




Salt stressed Spreading Orache leaf




1. <https://identify.plantnet.org/el/the-plant-list/species/Atriplex%20patula%20L./data>

Phytoremediation

- Using plants to clean up contaminated environments
- **Phytoextraction** = removal of contaminants through root system & stored in above ground shoots



Halophytes



- Plants adapted to grow in saline conditions
- **Accumulators** = take-up salt through roots and sequester in leaves and stem
 - Requires harvesting
- **Excretors** (recretohalophytes) = take-up salt through root system and translocate it to the above ground parts of the plant, then excrete the salts through specialized salt glands on the leaf surfaces

Halophytes



- Plants adapted to grow in saline conditions
- **Accumulators** = take-up salt through roots and sequester in leaves and stem
 - Requires harvesting
- **Excretors** (recretohalophytes) = take-up salt through root system and translocate it to the above ground parts of the plant, then excrete the salts through specialized salt glands on the leaf surfaces

Selected Species for Remediation



Side Oats Grama
Bouteloua curtipendula



Prairie Cordgrass
Sporobolus michauxianus



Switchgrass
Panicum virgatum



Sand Dropseed
Sporobolus cryptandrus

Excretors

Accumulators


Experimental Plots



Established background:
Cl⁻ levels = 20 mg/kg

In plot:
Cl⁻ levels = mean 365 mg/kg

Experimental Plots



Halophyte Type

- Excretor**
- Accumulator**

June 17, 2022



Side Oats
Grama

Sand
Dropseed

Switchgrass

Prairie
Cordgrass

Halophyte Type

Excretor
Accumulator



Prairie
Cordgrass

Switchgrass

Sand
Dropseed

Side Oats
Gramma

July 21st, 2022

August 18, 2022

Prairie
Cordgrass

Switchgrass

Side Oats Grama

Sand
Dropseed



End of Season



September 29th, 2022



Halophyte Type

Excretor

Accumulator

Composting - Switchgrass



September 30th, 2022



April 11th 2023

- 74% reduction in switchgrass biomass
- 60% reduction in side oats grama biomass

Regrowth of harvested material

June 1, 2023



Regrowth of Switchgrass



Un-harvested Switchgrass

June 22, 2023



Prairie Cordgrass

Switchgrass

Sand Dropseed

Side Oats Grama

Halophyte Type

Excretor

Accumulator

July 18, 2023

Side Oats Grama
Sand Dropseed
Prairie Cordgrass
Switchgrass

Halophyte Type

Excretor

Accumulator

August 28 2023



Prairie Cordgrass

Switchgrass

Side Oats Grama

Sand Dropseed

Halophyte Type

Excretor

Accumulator

Phytoremediation timelines (2023)



Species	Literature biomass production g DW /m ²	Observed biomass production g DW/m ²	Average Chloride uptake mg/m ²	Years for Remediation
Prairie Cordgrass				
Side Oats Grama				
Switchgrass				
Sand Dropseed				

Phytoremediation timelines (2023)



Species	Literature biomass production g DW /m ²	Observed biomass production g DW/m ²	Average Chloride uptake mg/m ²	Years for Remediation
Prairie Cordgrass	1000			
Side Oats Grama	1100			
Switchgrass	1500			
Sand Dropseed	25			

Phytoremediation timelines (2023)



Species	Literature biomass production g DW /m ²	Observed biomass production g DW/m ²	Average Chloride uptake mg/m ²	Years for Remediation
Prairie Cordgrass	1000	8904		
Side Oats Grama	1100	454.3		
Switchgrass	1500	7568		
Sand Dropseed	25	-		

Excretor


Accumulator

Phytoremediation timelines (2023)



Species	Literature biomass production g DW /m ²	Observed biomass production g DW/m ²	Average Chloride uptake mg/m ²	Years for Remediation
Prairie Cordgrass	1000	8904	41,052	
Side Oats Grama	1100	454.3	1,317	
Switchgrass	1500	7568	16,967	
Sand Dropseed	25	-	12,124	

Excretor



Accumulator



Phytoremediation timelines (2023)



Species	Literature biomass production g DW /m ²	Observed biomass production g DW/m ²	Average Chloride uptake mg/m ²	Years for Remediation
Excretor Prairie Cordgrass	1000	8904	41,052	1.9 years
	Side Oats Grama	1100	454.3	1,317
Accumulator Switchgrass	1500	7568	16,967	2.3 years
	Sand Dropseed	25	-	12,124 (2022)

Future research

- Analyze wet candle data for salt dispersion of excretor halophytes
- Mow parts of each plot to simulate roadside conditions
- Endophytes



Next steps



Thank you!

lauren.nawroth@queensu.ca

lyndsay.cartwright@trca.ca

zeeb-b@rmc.ca

www.trca.ca

